

SUPPLEMENTAL MATERIAL 1: FLORISTIC RELEVÉS AND VASCULAR CHECKLIST OF THE LIGURIAN 4090 HABITAT
BRYOPHYTES IN LIGURIAN 4090 HABITAT: ENDEMIC ORO-MEDITERRANEAN HEATHS WITH GORSE (*GENISTA DESOLEANA*)

Gabriele Berta^{1*}, Claudia Turcato², Giuseppina Barberis¹, Davide Dagnino¹

¹ Department of Earth, Environment and Life Sciences (DISTAV), Genoa University, Viale Benedetto XV, 16132 Genova, Italy.

²Ce.S.Bi.N. s.r.l., Via San Vincenzo 2 - 20th floor, 16121 Genova, Italy

*Correspondence: gabrieleberta95@gmail.com

Floristic relevés performed in three sites of Ligurian *Genista desoleana* hedgehog-heaths are here presented. Coverage values of vascular taxa were assigned as multiples of 5, with the exception of the 1% value; in case of coverages less than 1%, we used the "+" symbol. Coverage values of bryophytic taxa (Supplemental Material 3) were calculated using the following formula: $C = (N_i/N_{tot}) \times 100$ where N_i is the number of the occurrences of the species, and N_{tot} is the total number of occurrences of all the Bryophytes; then, we calculated the coverage value corresponding to each species contribution compared to the total coverage of the bryophytic layer.

Moreover, we provide the list of the vascular plants found in the three sites and their main ecological features. The nomenclature of taxa follows Bartolucci et al. (2018a, b, c, 2019a, b, 2020a, b) and Aleffi et al. (2020). Life form classification follows Raunkiær (1934): T scap = scapose therophytes, T par = parasitic therophytes, G bulb = bulbous geophytes, H bienn = biennial hemicryptophytes, H scap = scapose hemicryptophytes, H caesp = caespitose hemicryptophytes, Ch suffr = suffruticose chamaephytes, Ch frut = fruticose chamaephytes, NP = nanophanerophytes, P caesp = caespitose phanerophytes. The categories of humidity, temperature and light (i.e., Ucat, Tcat and Lcat, respectively) are estimated according to the Ellenberg indexes, obtained from Pignatti et al. (2017-2019), using the following thresholds: Ucat: 1-6 = dry, 7-12 = humid; Tcat: 1-6 = cryophytic, 7-12 = thermophytic; Lcat: 1-6 = shaded, 7-12 = illuminated. Life strategies are identified according to Grime (1977): S = Stress-tolerant, R = Ruderal, C = Competitive.

Site 1: Masso (Castiglione Chiavarese), 44.270109; 9.492761. Site 2: Monte Croce dei Tozzi, W side (Casarza Ligure), 44.263086; 9.467245. Site 3: Monte Croce dei Tozzi, W side (Casarza Ligure), 44.263400; 9.474050. Coordinates are expressed in WGS84 EPSG:4326.

SITE 1			
Performed by:	Turcato C., Dagnino D.		
Date:	24/04/2020		
Exposure	SE		
Altitude (meters a.s.l.)	295		
Acclivity (°):	15		
Vegetation type	<i>Genista desoleana</i> hedgehog-heath		
Height of tree vegetation (m):	-	Overall coverage (%):	40
Height of shrub vegetation (m):	0,3	Tree coverage (%):	-
Height of herbaceous vegetation (cm):	20	Shrub coverage (%):	30
Height of muscinal vegetation (cm):	1	Grass coverage (%):	10
Sampled area (mq):	5x5	Moss coverage (%):	1
Rocks (%):	-	Stones (%):	80
Species		Coverage (%)	
		bryophytic	herbaceous
<i>Genista desoleana</i> Vals.			30
<i>Cerastium ligusticum</i> Viv.			5
<i>Jasione montana</i> L.			1
<i>Agrostis castellana</i> Boiss. & Reut.			5
<i>Rumex acetosella</i> L.			1
<i>Saponaria ocymoides</i> L. subsp. <i>ocymoides</i>			+
<i>Bryum</i> sp.		0,4	
<i>Hypnum cupressiforme</i> Hedw. var. <i>cupressiforme</i>		0,27	
<i>Cephaloziella divaricata</i> (Sm.) Schiffn. var. <i>divaricata</i>		0,27	
<i>Dicranum scoparium</i> Hedw.		0,06	

SITE 2			
Performed by:	Turcato C., Dagnino D.		
Date:	24/04/2020		
Exposure	SO		
Altitude (meters a.s.l.)	224		
Acclivity (°):	40		
Vegetation type	<i>Genista desoleana</i> hedgehog-heath		
Height of tree vegetation (m):	-	Overall coverage (%):	30
Height of shrub vegetation (m):	0,4	Tree coverage (%):	-
Height of herbaceous vegetation (cm):	30	Shrub coverage (%):	30
Height of muscinal vegetation (cm):	0,5	Grass coverage (%):	1
Sampled area (mq):	5x5	Moss coverage (%):	1
Rocks (%):	30	Stones (%):	60
Coverage (%)			
Species		bryophytic	herbaceous
<i>Genista desoleana</i> Vals.			25
<i>Thymus vulgaris</i> L. subsp. <i>vulgaris</i>			5
<i>Peucedanum officinale</i> L.		+	
<i>Dianthus sylvestris</i> Wulfen		+	
<i>Festuca</i> sp.		1	
<i>Teucrium montanum</i> L.			1
<i>Trinia glauca</i> (L.) Dumort.		+	
<i>Buxus sempervirens</i> L.			+
<i>Dittrichia viscosa</i> (L.) Greuter subsp. <i>viscosa</i>		+	
<i>Juniperus oxycedrus</i> L.			+
<i>Euphorbia spinosa</i> L.			1
<i>Asperula</i> sp.		+	
<i>Weissia controversa</i> Hedw. var. <i>controversa</i>		0,917	
<i>Grimmia trichophylla</i> Grev.		0,083	

SITE 3			
Performed by:	Turcato C., Dagnino D.		
Date:	24/04/2020		
Exposure	SO		
Altitude (meters a.s.l.)	301		
Acclivity (°):	40		
Vegetation type	<i>Genista desoleana</i> hedgehog-heath		
Height of tree vegetation (m):	-	Overall coverage (%):	30
Height of shrub vegetation (m):	0,4	Tree coverage (%):	-
Height of herbaceous vegetation (cm):	50	Shrub coverage (%):	25
Height of muscinal vegetation (cm):	0,5	Grass coverage (%):	5
Sampled area (mq):	5x5	Moss coverage (%):	1
Rocks (%):	30	Stones (%):	60
Species		Coverage (%)	
		bryophytic	herbaceous
<i>Genista desoleana</i> Vals.			20
<i>Euphorbia spinosa</i> L.			5
<i>Thymus vulgaris</i> L. subsp. <i>vulgaris</i>			1
<i>Satureja montana</i> L. subsp. <i>montana</i>			5
<i>Centaurea aplolepa</i> Moretti subsp. <i>lunensis</i> (Fiori) Dostál		1	
<i>Linum trigynum</i> L.	+		
<i>Cuscuta</i> sp.	+		
<i>Festuca</i> sp.	5		
<i>Iberis umbellata</i> L.	+		
<i>Dianthus sylvestris</i> Wulfen	1		
<i>Teucrium montanum</i> L.			1
<i>Dittrichia viscosa</i> (L.) Greuter subsp. <i>viscosa</i>		+	
<i>Reichardia picroides</i> (L.) Roth	+		
<i>Asperula</i> sp.	+		
Liliacea	+		
Brassicacea	+		
<i>Weissia controversa</i> Hedw. var. <i>controversa</i>	0,75		
<i>Grimmia trichophylla</i> Grev.	0,125		
<i>Fissidens dubius</i> P.Beauv. var. <i>dubius</i>	0,125		

Taxon	Site 1	Site 2	Site 3	Life-form	Ucat	Tcat	Lcat	Life strategy
<i>Genista desoleana</i> Vals.	x	x	x	NP	dry	thermophytic	illuminated	S
<i>Festuca</i> sp.		x	x	H caesp	/	/	/	/
<i>Cuscuta</i> sp.			x	T par	/	/	/	R
Liliaceae			x	G bulb	/	/	/	/
Brassicaceae			x	/	/	/	/	/
<i>Cerastium ligusticum</i> Viv.	x			T scap	dry	thermophytic	illuminated	R
<i>Dittrichia viscosa</i> (L.) Greuter subsp. <i>viscosa</i>		x	x	H scap	dry	thermophytic	illuminated	R
<i>Iberis umbellata</i> L.			x	T scap	dry	thermophytic	illuminated	R
<i>Euphorbia spinosa</i> L.		x	x	Ch suffr	dry	cryophytic	illuminated	S
<i>Linum trigynum</i> L.			x	T scap	dry	thermophytic	illuminated	R
<i>Trinia glauca</i> (L.) Dumort.		x		H scap	dry	thermophytic	illuminated	S
<i>Juniperus oxycedrus</i> L.		x		P caesp	dry	thermophytic	illuminated	S-C
<i>Thymus vulgaris</i> L. subsp. <i>vulgaris</i>		x	x	Ch frut	dry	thermophytic	illuminated	S-C
<i>Dianthus sylvestris</i> Wulfen		x	x	H scap	dry	thermophytic	illuminated	S
<i>Teucrium montanum</i> L.		x	x	Ch suffr	dry	thermophytic	illuminated	S
<i>Agrostis castellana</i> Boiss. & Reut.	x			H caesp	dry	cryophytic	illuminated	C
<i>Satureja montana</i> L. subsp. <i>montana</i>			x	Ch suffr	dry	cryophytic	illuminated	S
<i>Rumex acetosella</i> L.	x			H scap	dry	cryophytic	illuminated	C
<i>Jasione montana</i> L.	x			H bienn	dry	thermophytic	illuminated	R
<i>Centaurea aplolepa</i> Moretti subsp. <i>lunensis</i> (Fiori) Dostál			x	H scap	dry	thermophytic	illuminated	S
<i>Reichardia picroides</i> (L.) Roth			x	H scap	dry	thermophytic	illuminated	S
<i>Asperula</i> cfr. <i>cynanchica</i> L. subsp. <i>cynanchica</i>		x	x	H scap	dry	thermophytic	illuminated	S
<i>Peucedanum officinale</i> L.		x		H scap	dry	cryophytic	illuminated	C-S
<i>Saponaria ocymoides</i> L. subsp. <i>ocymoides</i>	x			H scap	dry	cryophytic	illuminated	C-S
<i>Buxus sempervirens</i> L.		x		NP	dry	thermophytic	shaded	S-C

REFERENCES

- Aleffi M., Tacchi R. & Poponessi S., 2020. New checklist of the Bryophytes of Italy. *Cryptogamie, Bryologie* 41(13): 147-195. DOI: 10.5252/cryptogamie-bryologie2020v41a13
- Bartolucci F., Peruzzi L., Galasso G., Albano A., Alessandrini A., Ardenghi N. M. G., Astuti G., Bacchetta G., Ballelli S., Banfi E., Barberis G., Bernardo L., Bouvet D., Bovio M., Cecchi L., Di Pietro R., Domina G., Fascetti S., Fenu G., Festi F., Foggi B., Gallo L., Gottschlich G., Gubellini L., Iamonico D., Iberite M., Jiménez-Mejías P., Lattanzi E., Marchetti D., Martinetto E., Masin R. R., Medagli P., Passalacqua N. G., Peccenini S., Pennesi R., Pierini B., Poldini L., Prosser F., Raimondo F. M., Roma-Marzio F., Rosati L., Santangelo A., Scoppola A., Scortegagna S., Selvaggi A., Selvi F., Soldano A., Stinca A., Wagensommer R. P., Wilhalm T. & Conti F., 2018a. An updated checklist of the vascular flora native to Italy. *Plant Biosystems* 152(2): 179–303. DOI: 10.1080/11263504.2017.1419996
- Bartolucci F., Domina G., Ardenghi N. M. G., Banfi E., Bernardo L., Bonari G., Buccino G., Calvia G., Carruggio F., Cavallaro V., Chianese G., Conti F., Facioni L., Del Vico E., Di Gristina E., Falcinelli F., Forte L., Gargano D., Mantino F., Martino M., Mei G., Mereu G., Olivieri N., Passalacqua N. G., Pazienza G., Peruzzi L., Roma-Marzio F., Scafidi F., Scoppola A., Stinca A. & Nepi C., 2018b. Notulae to the Italian native vascular flora: 5. *Italian Botanist* 5: 71–81 DOI: 10.3897/italianbotanist.5.25892
- Bartolucci F., Domina G., Ardenghi N. M. G., Bacchetta G., Bernardo L., Buccino G., Buono S., Calderaro F., Calvia G., Carruggio F., Cavallaro V., Chianese G., Conti F., Facioni L., Del Vico E., Di Gristina E., Falcinelli F., Forte L., Gargano D., Mantino F., Martino M., Mei G., Mereu G., Olivieri N., Passalacqua N. G., Pazienza G., Peruzzi L., Roma-Marzio F., Scafidi F., Scoppola A., Stinca A. & Nepi C., 2018c. Notulae to the Italian native vascular flora: 6. *Italian Botanist* 6: 45–64. DOI: 10.3897/italianbotanist.6.30575
- Bartolucci F., Domina G., Alessandrini A., Angiolini C., Ardenghi N. M. G., Bacchetta G., Banfi E., Bolpagni R., Bonari G., Bräuchler C., Calvia G., Cancellieri L., Cannucci S., Carruggio F., Conti F., Cavallaro V., Fanfarillo E., Ferretti G., Festi F., Fiaschi T., Foggi B., Forte L., Fröhner S. E., Galasso G., Gestri G., Gottschlich G., Labadessa R., Lastrucci L., Lazzaro L., Mereu G., Morabito A., Mugnai M., Musarella C. M., Orsenigo S., Pazienza G., Pennesi R., Peruzzi L., Pierini B., Podda L., Prosser F., Rossi G., Scoppola A., Spampinato G., Stinca A., Tomaselli V., Zangari G. & Nepi C., 2019a. Notulae to the Italian native vascular flora: 7. *Italian Botanist* 7: 125–148. DOI: 10.3897/italianbotanist.7.36148
- Bartolucci F., Domina G., Ardenghi N. M. G., Bacaro G., Bacchetta G., Ballarin F., Banfi E., Barberis G., Beccarisi L., Bernardo L., Bonari G., Bonini F., Brullo S., Buono S., Buono V., Calbi M., Calderaro F., Calvia G., Cancellieri L., Cannavò S., Dagnino D., Esposito A., Fascetti S., Filibeck G., Fiorini G., Forte L., Galasso G., Gestri G., Gigante D., Gottschlich G., Gubellini L., Hofmann N., Lastrucci L., Lonati M., Lorenz R., Lunardi L., Magrini S., Mainetti A., Maiorca G., Mereu G., Messa Ballarin R. T., Minuto L., Mossini S., Musarella C. M., Nimis P. L., Passalacqua N. G., Peccenini S., Petriglia B., Podda L., Potenza G., Ravetto Enri S., Roma-Marzio F., Rosati L., Ruggero A., Spampinato G., Stinca A., Tiburtini M., Tietto C., Tomaselli V., Turcato C., Viciani D., Wagensommer R. P. & Nepi C., 2019b. Notulae to the Italian native vascular flora: 8. *Italian Botanist* 8: 95–116. DOI: 10.3897/italianbotanist.8.48626
- Bartolucci F., Domina G., Andreatta S., Angius R., Ardenghi N. M. G., Bacchetta G., Ballelli S., Banfi E., Barberis D., Barberis G., Bernardo L., Bertolli A., Bonari G., Bovio M., Briozzo I., Buccino G., Calvia G., Chianese G., Cibei C., Conti F., Copez M., Crisanti A., Dagnino D., Di Filippo A., Esposito A., Fanni S., Festi F., Forte L., Galasso G., Gentili R., Gottschlich G., Lattanzi E., Liguori P., Locci M. C., Longo D., Lonati M., Lucchese F., Marchetti D., Mariotti M. G., Menini F., Minuto L., Orrù G., Pala M. L., Passalacqua N. G., Pellegrino M., Pennesi R., Peruzzi L., Pinzani L., Pirastru G., Prosser F., Ravetto Enri S., Roma-Marzio F., Russo G., Scoppola A., Silletti G., Stinca A., Toffolo C., Tomaselli V., Tondi G., Trenchi M., Turcato C. & Nepi C., 2020a. Notulae to the Italian native vascular flora: 9. *Italian Botanist* 9: 71–86. DOI: 10.3897/italianbotanist.9.53429
- Bartolucci F., Domina G., Bagella S., Barberis G., Briozzo I., Calbi M., Caria M. C., Cavallaro V., Chianese G., Cibei C., Conti F., Dagnino D., Esposito A., Galasso G., Giacanelli V., Forte L., Gottschlich G., Lattanzi E., Longo D., Mei G., Merli M., Orsenigo S., Pau G. B., Pazienza G., Peccenini S., Pisani S., Rivieccio G., Roma-Marzio F., Scafidi F., Selvi F., Stinca A., Turcato C. & Nepi C., 2020b. Notulae to the Italian native vascularflora: 10. *Italian Botanist* 10:47–55 DOI: 10.3897/italianbotanist.10.60743
- Grime J. P., 1977. Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *The american naturalist* 111(982): 1169-1194. DOI: 10.1086/283244
- Pignatti S., Guarino R. & La Rosa M., 2017-19. Flora d'Italia (Edizione II). Edagricole - Edizioni Agricole di New Business Media srl, Bologna.
- Raunkiær C., 1934. The Life Forms of Plants and Statistical Plant Geography, being the collected papers of C. Raunkiær. Clarendon Press, Oxford.